arXiv:1603.07703v1 [math.CA] 24 Mar 2016

Kinks in the parametrically excited sine-Gordon equation and method of averaging

Vladimir Burd

Department of Mathematics, Yaroslavl State University, Russia

Abstract

Parametrically excited sine-Gordon equation is considered. Excitation is a fast oscillating periodic function with zero mean. Technique of classical method of averaging enables to construct the averaged equations in a variety of assumptions about driving amplitude. The averaged equation possesses kinks solutions. The results can be applied to the study of movement of Bloch walls for ferromagnetic crystals in the presence of a rapidly oscillating magnetic field and to describe the fluxon dynamics in long Josephson junctions.

Keywords

sine-Gordon equation, parametric excitation, method of averaging, kinks.

1 Introduction

As is well known, the sine-Gordon equation has a different physical applications. The sine-Gordon equation arose in the study of wave in quasi-one-dimensional ferromagnetic materials and the propagation of spin waves in superfluid phases A and B of helium ^3He. The sine-Gordon equation has been studied in connection with the Josephson junctions in the theory of superconductivity.

A significant number of studies have been devoted to development of technique of perturbation theory for systems close to integrable, in particular to systems close to the equation sine-Gordon [1–6]. The dynamics of kinks under the action of parametric perturbation was considered in [6,7,8]. This equation has a form

$$u_{tt}(x,t) - u_{xx}(t,x) + f\left(\frac{t}{\varepsilon}\right)\sin u(t,x) = 0. \tag{1}$$

Here $f(t)$ is a periodic function with zero mean and a constant amplitude, ε is small positive parameter. In the papers [7, 8] was found averaged equation. Averaged equation possesses π-kinks solutions.

The problem of constructing the averaged equation is solved in [7,8] as follows. It is introduced the Hamiltonian

$$H = \int\limits_{-\infty}^{\infty}\left(\frac{p^2}{2} + \frac{u_x^2}{2} - f\left(\frac{t}{\varepsilon}\right)\cos u\right)dx,$$

where $p = u_t$. Then, the series of canonical near-identical transformations is applied. It allows to remove the fast oscillating terms of lower orders from the Hamiltonian. Another

way of constructing the averaged dynamics was proposed in [6]. The solution parametrically excited sine-Cordon equation is sought in the form of a Fourier series with slowly varying coefficients on the time scale ω^{-1}, ω being the frequency of the rapidly varying perturbation. .

In this paper we propose a third method of constructing of averaged equation for (1). It is the classical method of averaging (see [9,10]). Our approach is similar to the method that was used in the study of the stability of the upper equilibrium of the pendulum with a vertically vibrating pivot [11].

2 Constructing of averaged equations

We denote by $f_{-1}(t)$ the periodic function with zero mean value whose derivative $f'_{-1}(t)$ satisfies the equality

$$f'_{-1}(t) = f(t).$$

From equation (1) we move to equivalent system

$$\begin{array}{l} u_t = p - \varepsilon f_{-1}\left(\frac{t}{\varepsilon}\right)\sin u \\ p_t = u_{xx} + \varepsilon f_{-1}\left(\frac{t}{\varepsilon}\right) p\cos u - \frac{1}{2}\varepsilon^2\left(f_{-1}\left(\frac{t}{\varepsilon}\right)\right)^2 \sin 2u. \end{array} \tag{2}$$

This transition is essentially a transition from a recording of system in the form of Lagrange to recording of system in the Hamiltonian form. In system (2) turn to fast time $\tau = \frac{t}{\varepsilon}$. We obtain a system

$$\begin{array}{l} u_\tau = \varepsilon p - \varepsilon^2 f_{-1}(\tau)\sin u \\ p_\tau = \varepsilon u_{xx} + \varepsilon^2 f_{-1}(\tau) p\cos u - \frac{1}{2}\varepsilon^3 (f_{-1}(\tau))^2 \sin 2u. \end{array} \tag{3}$$

The right-hand sides of the system (3) are proportional to small parameter ε. This is a standard form for applying the method of averaging. We make regular substitution of method of averaging

$$\begin{array}{l} u = \xi + \varepsilon^2 v_2(\tau,\xi) + \varepsilon^3 v_3(\tau,\xi), \\ p = \eta + \varepsilon^2 w_2(\tau,\xi,\eta) + \varepsilon^3 w_3(\tau,\xi,\eta). \end{array} \tag{4}$$

The change should eliminate a variable τ from the right-hand side of the system (3) up to terms of the fourth order. Therefore we need to get the system

$$\begin{array}{l} \xi_\tau = \varepsilon\eta + \varepsilon^2 A_2(\xi,\eta) + \varepsilon^3 A_3(\xi,\eta) + O(\varepsilon^4), \\ \eta_\tau = \varepsilon\xi_{xx} + \varepsilon^2 B_2(\xi,\eta) + \varepsilon^3 B_3(\xi,\eta) + O(\varepsilon^4). \end{array} \tag{5}.$$

Let us find coefficients A_2, A_3, B_2, B_3 of system (5). By substituting (4) into (3) and replacing ξ_τ and η_τ with right-hand sides of the system (5) we get

$$\begin{array}{l} \varepsilon\eta + \varepsilon^2 A_2(\xi,\eta) + \varepsilon^3 A_3(\xi,\eta) + \varepsilon^2 v_{2\tau}(\tau,\xi) + \varepsilon^3 v_{2\xi}(\tau,\xi)\eta + \varepsilon^3 v_{3\tau}(\tau,\xi) + O(\varepsilon^4) = \\ \varepsilon\eta - \varepsilon^2 f_{-1}(\tau)\sin\xi + \varepsilon^3 w_2(\tau,\xi,\eta) + O(\varepsilon^4), \\ \varepsilon\xi_{xx} + \varepsilon^2 B_2(\xi,\eta) + \varepsilon^3 B_3(\xi,\eta) + \varepsilon^2 w_{2\tau}(\tau,\xi,\eta) + \\ \varepsilon^3 w_{2\xi}(\tau,\xi,\eta)\eta + \varepsilon^3 w_{2\eta}(\tau,\xi,\eta)\xi_{xx} + \varepsilon^3 w_{3\tau}(\tau,\xi,\eta) + O(\varepsilon^4) \\ = \varepsilon\xi_{xx} + \varepsilon^3\frac{\partial^2}{\partial x^2}(v_2(\tau,\xi)) + \varepsilon^2 f_{-1}(\tau)\eta\cos\xi - \varepsilon^3\frac{1}{2}(f_{-1}(\tau))^2\sin 2\xi + O(\varepsilon^4). \end{array} \tag{6}$$

Equating the coefficients of power of ε^2 we obtain

$$v_{2\tau}(\tau,\xi)+A_2(\xi,\eta)=-f_{-1}(\tau)\sin\xi,\quad w_{2\tau}(\tau,\xi,\eta)+B_2(\xi,\eta)=f_{-1}(\tau)\eta\cos\xi.$$

The function $A_2(\xi,\eta)$ is defined as the mean value on τ of the function $-f_{-1}(\tau)\sin\xi$. Hence we obtain $A_2(\xi,\eta)\equiv 0$. Then the function $v_2(\tau,\xi)$ is uniquely determined as periodic function on τ with zero mean value. The function is defined by formula $v_2(\tau,\xi)=-(\int f_{-1}(s)ds)\sin\xi$. Similarly we find that $B_2(\xi,\eta)\equiv 0$. The function $w_2(\tau,\xi,\eta)$ is periodic function on τ with zero mean value and $w_2(\tau,\xi,\eta)=(\int f_{-1}(s)ds)\eta\cos\xi$. Equating coefficients of power ε^3 in (6) we obtain

$$A_3(\tau,\xi)+v_{3\tau}(\tau,\xi)+v_{2\xi}(\tau,\xi)\eta+v_{2\eta}(\tau,\xi)\xi_{xx}=w_2(\tau,\xi,\eta),$$

$$B_3(\xi,\eta)+w_{3\tau}(\tau,\xi,\eta)+w_{2\xi}(\tau,\xi,\eta)\eta+w_{2\eta}(\tau,\xi,\eta)\xi_{xx}=-\frac{1}{2}(f_{-1}(\tau))^2\sin 2\xi+\frac{\partial^2}{\partial x^2}(v_2(\tau,\xi)).$$

Mean values of functions $v_{2\xi}(\tau,\xi)\eta$, $v_{2\eta}(\tau,\xi)\xi_{xx}$, $w_2(\tau,\xi,\eta)$ are zero. Then $A_3(\xi,\eta)\equiv 0$. The same arguments shows that $B_3(\tau,\xi,\eta)$ is determined by formula

$$B_3(\xi,\eta)=\frac{1}{2}\langle(f_{-1}(\tau))^2\rangle\sin 2\xi,$$

where $\langle f_{-1}^2(\tau)\rangle$ is mean value of periodic function $f_{-1}^2(\tau)$. Therefore the averaged system has the form

$$\begin{aligned}&\xi_\tau=\varepsilon\eta,\\&\eta_\tau=\varepsilon\xi_{xx}-\varepsilon^3\tfrac{1}{2}\langle(f_{-1}(\tau))^2\rangle\sin 2\xi.\end{aligned}$$

Going back to the time $t=\varepsilon\tau$ we obtain the system

$$\begin{aligned}&\xi_t=\eta,\\&\eta_t=\xi_{xx}-\varepsilon^2\tfrac{\Delta}{2}\sin 2\xi,\end{aligned}$$

where

$$\Delta=\langle(f_{-1}(\tau))^2\rangle.$$

This system can be written as an equation of the second-order

$$\xi_{tt}-\xi_{xx}+\varepsilon^2\frac{\Delta}{2}\sin 2\xi=0. \tag{7}$$

The equation (7) has the solutions

$$u(x,t)=2\arctan\left[\exp\left((x-ct)\frac{\varepsilon\sqrt{\Delta}}{\sqrt{1-c^2}}+\delta\right)\right],$$

where δ is constant. These solutions are π kinks.

It was assumed above that the function $f(t/\varepsilon)$ has a constant amplitude. Now we assume that amplitude of excitation have the order of $1/\varepsilon$. Let excitation is

$$\frac{1}{\varepsilon}f\left(\frac{t}{\varepsilon}\right).$$

Parametric excitation of the sine-Gordon equation now has the form

$$u_{tt}(x,t) - u_{xx}(t,x) + \frac{1}{\varepsilon} f\left(\frac{t}{\varepsilon}\right) \sin u(t,x) = 0. \tag{8}$$

We again denote by $f_{-1}(t)$ the periodic function with zero mean value that derivative $f'_{-1}(t)$ satisfies the equality

$$f'_{-1}(t) = f(t).$$

From equation (8) we pass to the equivalent system of equations

$$\begin{aligned} u_t &= p - f_{-1}\left(\tfrac{t}{\varepsilon}\right) \sin u \\ p_t &= u_{xx} + f_{-1}\left(\tfrac{t}{\varepsilon}\right) p \cos u - \tfrac{1}{2}\left(f_{-1}\left(\tfrac{t}{\varepsilon}\right)\right)^2 \sin 2u. \end{aligned}$$

Then move on to the fast time $\tau = t/\varepsilon$. We obtain the system

$$\begin{aligned} u_\tau &= \varepsilon(p - f_{-1}(\tau) \sin u) \\ p_\tau &= \varepsilon(u_{xx} + f_{-1}(\tau) p \cos u - \tfrac{1}{2}(f_{-1}(\tau))^2 \sin 2u). \end{aligned}$$

The averaged system is defined in a first approximation and has the form

$$\begin{aligned} \xi_\tau &= \varepsilon\eta \\ \eta_\tau &= \varepsilon(\xi_{xx} - \tfrac{1}{2}\langle (f_{-1}(\tau))^2 \rangle \sin 2\xi). \end{aligned}$$

We assume $\Delta = \langle f_1(\tau)^2 \rangle$ and record the average system in the form of a second-order equation in time t

$$\xi_{tt} - \xi_{xx} + \frac{\Delta}{2} \sin 2\xi = 0. \tag{9}.$$

The equation (9) has the π- kinks

$$u(x,t) = 2 \arctan\left[\exp\left((x - ct)\frac{\sqrt{\Delta}}{\sqrt{1-c^2}} + \delta\right)\right], \quad \delta = const.$$

Let's look at the excitement of the following form

$$u_{tt}(x,t) - u_{xx}(t,x) + \left(1 + \frac{1}{\varepsilon} f\left(\frac{t}{\varepsilon}\right)\right) \sin u(t,x) = 0. \tag{10}$$

Note that in the case of one variable, equation (10) is transformed into the equation of motion of the pendulum with a vertically oscillating pivot and small amplitude. In the notations introduced earlier system of equations is equivalent to the equation (10) has the form

$$\begin{aligned} u_t &= p - f_{-1}\left(\tfrac{t}{\varepsilon}\right) \sin u, \\ p_t &= u_{xx} - \sin u + f_{-1}\left(\tfrac{t}{\varepsilon}\right) p \cos u - \tfrac{1}{2}\left(f_{-1}\left(\tfrac{t}{\varepsilon}\right)\right)^2 \sin 2u. \end{aligned} \tag{11}$$

We go to a fast time $\tau = t/\varepsilon$. Then we average system (11). Averaged system of the first approximation in time t can be written as equation of second order

$$\xi_{tt} - \xi_{xx} + \sin\xi + \frac{\Delta}{2} \sin 2\xi = 0. \tag{12}$$

where $\Delta = \langle f_1(\tau)^2 \rangle$. Thus averaged equation is the double sine-Gordon equation. The kinks solution of equation (12) may be written in the form

$$\xi(t,x) = 2\arctan\left[\frac{1}{\sqrt{1+\Delta/2}} csch\left(\sqrt{1+\Delta/2}\frac{x-ct}{\sqrt{1-c^2}}\right)\right].$$

Similarly, the following parametric perturbed equation

$$u_{tt} - u_{xx} + \left(1 + f\left(\frac{t}{\varepsilon}\right)\right)\sin u = 0. \tag{13}$$

is considered. The equivalent system of equations is

$$\begin{aligned} u_t &= p - \varepsilon f_{-1}\left(\tfrac{t}{\varepsilon}\right)\sin u \\ p_t &= u_{xx} - \sin u + \varepsilon f_{-1}\left(\tfrac{t}{\varepsilon}\right) p\cos u - \tfrac{1}{2}\varepsilon^2\left(f_{-1}\left(\tfrac{t}{\varepsilon}\right)\right)^2 \sin 2u. \end{aligned}$$

The averaged equation for (13) has the form

$$\xi_{xx} - \xi_{xx} + \sin\xi + \varepsilon^2\frac{\Delta}{2}\sin 2\xi = 0.$$

3 Conclusion

The classical method of averaging is proposed for the construction of the averaged dynamics in parametrically perturbed sine-Gordon equation. The essential point is transition to an equivalent system of first order equations. The averaging method is also applicable to the case of the direct driving force in the sine-Gordon equation and driven damped sine-Gordon model. This method can be used to analyze the dynamics of kinks on rotationg and oscillating background.

References

[1] Yu.S. Kivshar, B.A. Malomed Dynamics of solitons in nearly by integrable systems, Reviews of Modern Physics, 1989, vol. 61, no. 4, pp. 763–915.

[2] J.C. Fernandez, Th. Passot, H. Politano, G. Reinisch, M. Taki Sturm-Liouville description of sine-Gordon soliton dynamics, Physical Review B, 1988, vol. 37, no. 13, pp. 7342–7347.

[3] R. Grauer, Yu. S. Kivshar Chaotic and phase-locked breather dynamics in the damped and parametrically driven sine-Gordon equation, Physical Review B, 1993, vol. 48, no. 6, pp. 4791–4800.

[4] S. Sakai, M.R. Samuelsen, O.H. Olsen Perturbation analysis of a parametrically changed sine-Gordon equation, Physical Review B, 1987, vol. 36, no. 1, pp. 217–225.

[5] Yu.S. Kivshar, N. Gr∅nbech-Jensen, M.R. Samuelsen, π kinks in a parametricaly driven sine-Gordon chain, Physical Review B, 1992, vol. 45, no. 14, pp. 7789–7794.

[6] Yu. S. Kivshar, N. Gr∅nbech-Jensen, R.D. Parmentier Kinks in the presence of rapidly varying perturbations, Physical Review E, 1994, vol. 49, no. 5, pp. 4542–4551.

[7] V. Zharnitsky, I. Mitkov, M. Levi Parametrically forced sine-Gordon equation and domain wall dynamics in ferromagnets, Physical Review B, 1998, vol. 57, no. 9, pp. 5033–5035.

[8] I. Mitkov, V.Zharnitsky π-Kinks in the parametrically driven sine-Gordon equation and applications, Physica D, 1998, vol. 123, pp. 301–307.

[9] N.N. Bogoliubov and Yu.A. Mitropolskiy Asimptotic methods in the theory of nonlinear oscillatios. Gordon and Breach, New York, 1961.

[10] J. Sanders, F. Verhulst, J. Murdoc Averaging methods in nonlinear dynamical systems, 2ed., Springer, Applied Mathematical Sciences, vol. 59, Springer, 2007,

[11] V. Burd V. Method of averaging for differential equations on an infinite interval. Theory and applications, Lecture Notes in Pure and Applied Mathematics, vol. 255. Chapman&Hall/CRC, Boca Raton, London, New York, 2007.